



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED
LAWRENCE
BERKELEY LABORATORY

MAR 14 1984

LIBRARY AND
DOCUMENTS SECTION

Submitted to Physical Review Letters


MEASUREMENTS OF HIGH p_T LIGHT FRAGMENTS AT C.M. 90°
IN 800 A·MeV C + C COLLISIONS

H. Hamagaki, X.X. Bai, O. Hashimoto, S. Kadota,
K. Kimura, Y. Miake, S. Nagamiya, K. Omata,
Y. Shida, I. Tanihata, and N. Yoshikawa

January 1984

For Reference

Not to be taken from this room



LBL-17276
e.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

**Measurements of High p_T Light Fragments at C.M. 90°
in 800 A-MeV C + C Collisions**

H. Hamagaki, X. X. Bai^{a)}, O. Hashimoto, S. Kadota,
K. Kimura^{b)}, Y. Miake^{c)}, S. Nagamiya^{c)}, K. Omata,
Y. Shida, I. Tanihata, and N. Yoshikawa

Institute for Nuclear Study, University of Tokyo,
Tanashi-city, Tokyo 188, Japan

and

Nuclear Science Division, Lawrence Berkeley Laboratory,
University of California, Berkeley, California 94720

Abstract

The spectra of high- p_T fragments, p , d , and t , have been measured at c.m. 90° in 800 A-MeV C + C collisions for p_T up to 1.8 GeV/c (proton), 2.5 GeV/c (deuteron) and 2.8 GeV/c (triton), using the Heavy Ion Spectrometer System (HISS) at the Bevalac. Invariant cross sections were measured down to eight orders of magnitude lower than those for production of low momentum protons. The proton spectrum shows a significant deviation from the Boltzmann shape in the high-energy region. The spectra of deuterons and tritons are explained well by, respectively, squaring and cubing the observed proton spectra.

PACS Numbers: 25.70Fg, 25.70Bc.

In high-energy nucleus-nucleus collisions it is important to study particle emission in the high- p_T region for various reasons. If the projectile and target consist of m - and n -nucleons, respectively, the maximum kinetic energy, E_{\max} , for protons produced at 90° in the nucleon-nucleon (NN) c.m., center of mass, frame is given by

$$E_{\max} = (\gamma-1)m_p c^2 + \left[\frac{2mn}{m+n} - 1 \right] \gamma \beta^2 m_p c^2 \quad (1)$$

where γ and β represent the familiar Lorentz transformation variables that define the NN c.m. frame relative to the laboratory frame. At $E_{\text{Beam}}^{\text{Lab}} = 800 \text{ A} \cdot \text{MeV}$, $E_{\max} = 182 \text{ MeV}$ for $m = n = 1$ (free NN collisions) and $E_{\max} = 517 \text{ MeV}$ for $m = 1$ and $n \rightarrow \text{large}$ (proton-nucleus collisions), whereas E_{\max} reaches 3.9 GeV for $\text{C} + \text{C}$ collisions ($m = n = 12$). Therefore, emission of protons at c.m. 90° having kinetic energies above 517 MeV is possible only in nucleus-nucleus collision.

We may think of several mechanisms that induce proton emission in the high- p_T region. The first one is random multiple NN collisions. Roughly speaking, if an average momentum transfer per NN collision is Δp , then the total momentum transfer after the n -th NN collision would reach $\sim \sqrt{n} \Delta p$. Therefore, if n is sufficiently large, emission of high- p_T protons is possible. The second mechanism is attributable to cooperative processes of several nucleons. A local hot and dense region might be created in the nucleus-nucleus collision through a macroscopic compression of matter. Then, nucleons might be strongly correlated which may induce high-momentum transfers on the order of $\hbar/\Delta x$, where Δx is a typical correlation length. In the first process, a Boltzmann-type spectrum is expected in the low- p_T region, whereas particle emission would be suppressed relative to that in the high- p_T region because of the limited number of NN collisions. On the other hand, in the second process a new exponential component may appear in the high- p_T region.

In the past, the proton energy distribution at c.m. 90° in $800 \text{ A} \cdot \text{MeV} \text{ C} + \text{C}$ collisions has been measured for proton energies up to 700 MeV .¹ The spectrum shows a "shoulder-arm" shape with the turning point between the shoulder and arm located at $E_p^{\text{c.m.}} \sim 200 \text{ MeV}$. Above this energy the spectrum shape approaches an exponential,

$$E(d^3\sigma/dp^3) \propto \exp(-E_p^{\text{c.m.}}/E_0), \quad (2)$$

with $E_0 \approx 68$ MeV. In the present experiment we have extended these measurements to higher proton energies, where the cross section is four orders of magnitude lower than that at $E_p^{\text{c.m.}} = 700$ MeV, to study if other new features start to appear.

A layout of the experimental setup is shown in Fig. 1. A two-arm spectrometer system was constructed upon a base of the Heavy Ion Spectrometer System² (HISS) at the Bevalac. A magnetic field of 17.5 kG was applied over a region of 2-meter diam. \times 1-meter gap. A carbon target was mounted inside the magnet gap. The magnetic field served two purposes: to eliminate low-energy particles from the event trigger and to analyze momentum of high-energy particles. Here, we report measurements of inclusive spectra by the A-arm shown in Fig. 1. Two sets of three-plane multi-wire proportional chambers (WCA1; 20 cm \times 20 cm, WCA2; 20 cm \times 30 cm), two sets of six-plane drift chambers (DC1, DC2; 1 m \times 2 m), and 47 sets of plastic scintillators (TA1, TA2, TA3) were used to measure particle tracks, dE/dx , and time-of-flight. Particle identification was almost perfect up to $p_{\text{Lab}} \approx 4$ GeV/c. The momentum resolution was typically 6 % at $p_{\text{Lab}} = 3$ GeV/c, and the absolute cross section value is reliable to within ± 20 %. Details of the spectrometer performance will be discussed elsewhere.

Fig. 2 shows the observed proton energy distributions at c.m. 90° in 800 A-MeV C + C. Open circles indicate the present results, whereas closed circles show the data reported in Ref. 1. The open and closed circles have no kinematical overlaps. We have checked the consistency of the present data for protons, deuterons, and tritons with those reported in Ref. 1 at 45° in the laboratory frame where two data sets overlap.

These data cover invariant cross sections over eight orders of magnitudes. The solid line in Fig. 1 shows our previous fit¹ with an exponential shape given by Eq. (2). Clearly, the data show a smooth inward deviation from this exponential shape not only in the low-energy region below about 200 MeV but also in the high-energy region above 700 MeV. It should be noted that the slope of the spectrum at the high energy region is quite similar to that of negative pions.¹ No second slow exponential components have been observed at the level of cross sections down to $\approx 10^{-8}$ of the major cross sections.

Thus far, the c.m. 90° spectra have been studied extensively for the purpose of studying the participant dynamics, since at this angle the influence from the projectile and target fragments would be the smallest. It has been known that any thermal model³⁻⁵ cannot explain the data, because they predict a pure exponential shape over the entire kinematic domain of emitted protons. We, therefore, compared our data with (a) a generalized thermal model called the phase-space model^{6,7} and (b) the cascade model using the code of Cugnon⁸. As shown in Fig.2, the phase-space model seems consistent with the data. Although it slightly overestimates the yield in the low-energy region, it explains inward deviations from the exponential shape in both low- and high-energy regions. In regard to the cascade model, calculations were possible only up to $E_p^{c.m.} \approx 850$ MeV because of the limitation of the computation time. In the low-energy region, the model gives a reasonable fit to the data, but it gives a steeper slope than the data in the high-energy region, and at $E_p^{c.m.} \approx 850$ MeV the calculated cross section is by a factor of 6 smaller than the observed one. These comparisons tell us that the observed inward deviation from the exponential in the high- p_T region is likely due to a limited phase space available in this region, but the microscopic mechanisms that limit the available phase space are not totally due to the limited number of the random multiple NN collisions. In other words, some coherent mechanism other than random multiple NN collisions might be needed to fill up the available phase space.

It may be worthwhile to note that a modified thermal model proposed by Siemens and Rasmussen,⁹ that is, the radial explosion model in which an exploding flow is superposed on the thermal spectrum, may also be consistent with the data at least in a qualitative manner. In this model, the apparent temperature is given by $T_{app} = T\gamma_0^{-1}(1-\beta_0/\beta)^{-1}$, where T is the temperature of the system, $\beta_0 c$ is the flow velocity, βc is the velocity of the emitted particles, and $\gamma_0 = (1-\beta_0^2)^{-1/2}$. The value of T_{app} , thus, decreases as β increases and it approaches constant as β approaches 1. This is consistent with the observed smooth variation of the slope for proton cross sections and with the fact that the slope of the spectrum at high-energy region is similar to that for pions. Quantitatively, however, no detailed comparison is currently available.

Shown in Fig.3 are the deuteron and triton spectra at c.m. 90° from the same collision. Deuteron cross sections cover about 7 orders of magnitudes and triton about 5 orders. In both cases we have tried to compare the data with squares and cubes of the observed proton cross sections, respectively, as expected from a coalescence model.^{10,11} Amazingly, for over the six orders of magnitudes the cross sections satisfy the coalescence relation of

$$E_A (d^3 \sigma / dp_A^3) = C_A [E_p (d^3 \sigma / dp_p^3)]^A \quad \text{for } \vec{p}_A = A \cdot \vec{p}_p, \quad (3)$$

with $C_A = (3.5 \pm 0.7) \times 10^{-5} [(\text{mb}/\text{sr})\text{c}^3 / (\text{GeV})^2]^{-1}$ for d and $= (1.0 \pm 0.3) \times 10^{-9} [(\text{mb}/\text{sr})\text{c}^3 / (\text{GeV})^2]^{-2}$ for t . These values are consistent with the values reported in Ref.1 to within the quoted errors.

We also note that both the deuteron and triton spectra are not of an exponential type, reflecting the fact that the proton spectrum is not exponential. Therefore, from Fig.3 we conclude that the light composite particles are formed through final state interactions among nucleons over a wide range of kinetic energy.

In summary, we have observed proton cross sections at c.m. 90° in 800 A·MeV C + C collisions over 8 orders of magnitudes in cross section. A substantial inward deviation from the exponential shape is observed not only in the low-energy region below 200 MeV but also in the high-energy region above 700 MeV. The inward deviation at high energy is most likely due to the decreased phase space for the emission of high-energy protons, but the microscopic mechanism to fill up the phase space seems not totally due to the random NN collisions. We have also measured deuteron and triton spectra in the same conditions and found that the power law [Eq.(3)] holds extremely well between proton spectrum and the spectra of composite particles, even 7 orders of magnitudes in the cross sections. This result implies that the final state interactions are important in the formation of these composite particles in wide ranges of kinetic energies. Finally, we mention that experimental research to clarify microscopic mechanism of proton emission in the high- p_T region is currently in progress in our group by measuring two-proton correlations using the B-arm spectrometer show in Fig.1.

We would like to thank the members of the HISS group for collaboration at various stages in the experiment, and R. Fuzesy and J. Bistirlich for their technical assistance. Thanks also go to J. Knoll for his calculation of the phase-space model. We also thank H. Steiner, S. Schnetzer, Y. Kitazoe, and O. Chamberlain for stimulating discussions and advises. This work was supported by the Director, Division of Nuclear Physics of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and by the Collaboration Program between Institute for Nuclear Study (University of Tokyo) and Lawrence Berkeley Laboratory.

References

- a) Present address: Institute of Atomic Energy, P. O. Box 275, Beijing, People's Republic China.
 - b) Present address: Department of Physics, Kyushu University, Hakozaki, Higashi-ku, Fukuoka-city, Fukuoka 812, Japan.
 - c) Present address: Department of Physics, University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan.
1. S. Nagamiya, M.-C. Lemaire, E. Moeller, S. Schnetzer, G. Shapiro, H. Steiner, and I. Tanihata, *Phys. Rev. C* 24, 971 (1981).
 2. D. E. Greiner, *Nucl. Phys.* A400, 325c (1983).
 3. G. D. Westfall, J. Gosset, P. J. Johansen, A. M. Poskanzer, W. G. Meyer, H. H. Gutbrod, A. Sandoval, and R. Stock, *Phys. Rev. Lett.* 37, 1202 (1976).
 4. J. Gosset, J. I. Kapusta, and G. D. Westfall, *Phys. Rev. C* 18, 844 (1978).
 5. S. Das Gupta and A. Z. Mekjian, *Phys. Rep.* C72, 131 (1981).
 6. J. Knoll, *Phys. Rev. C* 20, 773 (1979).
 7. The calculation by the phase-space model was done by J. Knoll, private communication (1983).
 8. J. Cugnon, T. Mizutani, and J. Vandermeulen, *Nucl. Phys.* A352, 505 (1981).
 9. P. J. Siemens and J. O. Rasmussen, *Phys. Rev. Lett.* 42, 880 (1979).
 10. H. H. Gutbrod, A. Sandoval, P. J. Johansen, A. M. Poskanzer, J. Gosset, W. G. Meyer, G. D. Westfall, and R. Stock, *Phys. Rev. Lett.* 37, 667 (1976).
 11. M.-C. Lemaire, S. Nagamiya, S. Schnetzer, H. Steiner, and I. Tanihata, *Phys. Lett.* 85B, 38 (1979).

Figure Captions

FIG. 1 Layout of the experimental setup.

FIG. 2 Proton energy distribution measured at c.m. 90° in 800 A-MeV C + C collisions. Data are compared with the phase-space model (dot-dashed curve) and the cascade model (dashed curve).

FIG. 3 Deuteron and triton spectra measured at c.m. 90° in 800 A-MeV C + C collisions, as compared with squares and cubes, respectively, of the observed proton spectrum.






Fig. 1



XBL 839-872A

Fig. 2



XBL839-871

Fig. 3

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720