LBL-19305
Preprint c.2

Lawrence Berkeley Laboratory
UNIVERSITY OF CALIFORNIA

RECEIVED
LAWRENCE
BERKELEY LABORATORY

MAY 16 1985

LIBRARY AND
DOCUMENTS SECTION

Submitted to Physics Letters, B

CONVERGENCE OF PERTURBATION SERIES
IN THE MICROCANONICAL FORMULATION
OF QUANTUM FIELD THEORIES

A. Iwazaki

March 1985

TWO-WEEK LOAN COPY
This is a Library Circulating Copy
which may be borrowed for two weeks.

Prepared for the U.S. Department of Energy under Contract DE-AC03-76SF00098

LBL-19305
c.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

LBL-19305

Convergence of Perturbation Series in the Microcanonical Formulation of Quantum Field Theories

Aiichi Iwazaki

Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

This work was supported in part by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

LBL-19305

Convergence of Perturbation Series in the Microcanonical Formulation of Quantum Field Theories

Aiichi Iwazaki

Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Abstract

It is shown that the perturbation series in the microcanonical formulation of scalor $\lambda\phi^4$ and gauge theories on the lattice is absolutely convergent for small enough expansion parameter, λ. These series can be either the weak (λ) or the strong (λ^{-1}) coupling expansion series.

This work was supported in part by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

Recently, the microcanonical quantization method[1] has received much attention as a means of performing numerical calculations. Notably, it has been used in lattice gauge theories. Since the method is quite new to particle physicists, it is important to examine in detail the basis[2] of this quantum formulation.

We shall show in this letter that for both the strong and weak coupling expansion of scalar field theories or gauge theories, the perturbation series are convergent if we quantize in a finite volume with the microcanonical ensemble density.[1] It is possible, in principle, to sum these perturbation series.

As is well known,[3] the perturbation series in the standard formulations (Feynman's path integral or canonical quantization) is not convergent, even if we use an ultraviolet cut off and a finite volume. Although the series in the standard perturbation expansion can be summed up in some cases with appropriate methods of summation (Borel summation, etc.), the final results may still depend on the methods used. Furthermore, so called non-perturbative effects can not be recovered unambiguously from the standard perturbation expansion. On the other hand, the convergence of the microcanonical perturbation series implies that we can obtain all the effects. The underlying reason for the convergence of the series is that the microcanonical density is expanded with respect to an interaction Hamiltonian, but not with respect to the total energy, which in general depends itself on an expansion parameter and should be determined self-consistently (see below).

First, let us recapitulate briefly the microcanonical quantization method.[1,2] Consider a scalar theory on a d-dimensional lattice. The action is

$$S \equiv S_0 + \lambda S_I \quad , \quad S_0 \equiv \frac{V}{N}\left\{\sum_{x,\mu} \frac{(\phi_{x+\mu} - \phi_x)^2}{2a^2} + \sum_{\mu} \frac{m^2}{2} \phi_x^2\right\}$$

and

$$S_I = \sum_x \frac{V}{N} \phi_x^4 \quad , \tag{1}$$

where a, m and λ are the lattice spacing, bare mass and bare coupling constant, respectively, and where x is the vector belonging to a lattice point and μ is a unit vector in the μ direction. The volume $V = Na^d$ is taken to be finite. According to the microcanonical method, we compute the reguralized Green's function as follows

$$G(x_1 \ldots x_n) = \lim_{N\to\infty} \frac{1}{Z} \int d\mu\, \phi(x_1) \ldots \phi(x_n)\delta(E - H)$$

$$Z \equiv \int d\mu\, \delta(E - H) \quad , \tag{2}$$

with $H = \sum_{x=1}^{N} \frac{P_x^2}{2} + S$ and $E = \frac{N}{2} + \langle S\rangle$.

The measured used in (2) is given by

$$d\mu \equiv \prod_{x=1}^{N} dP_x d\phi_x \tag{3}$$

and the average $\langle S\rangle$ in E is

$$\langle S\rangle = \frac{\int \prod_{x=1}^{N} d\phi_x e^{-S} S}{\int \prod_{x=1}^{N} d\phi_x e^{-S}} \quad . \tag{4}$$

We shall show below that the Green's function in (2) is identical to the one given in the standard formulation. In a previous paper,[2] we have shown this equivalence perturbatively. It is easy to see that the formula in (2) is the same as the one used in the numerical calculations of Ref. 4. We note that the quantity <S> may also be obtained as

$$\lim_{N\to\infty} \langle S\rangle/N = \lim_{N\to\infty} \frac{1}{Z} \int d\mu\, S\delta(E - H)/N \quad . \tag{5}$$

This is a consistency condition, by use of which the total energy E (or <S>) is to be evaluated.

Now we turn to the microcanonical perturbation theory and show the convergence of its series. The perturbation series we consider is defined by expanding the microcanonical density with respect to λ,

$$\delta(E - H) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \frac{d^n}{dE^n} \delta(E - H_0)(\lambda S_I)^n \tag{6}$$

$$\text{with} \quad H = \sum_{\chi=1}^{N} \frac{p_\chi^2}{2} + S_0 \quad .$$

To demonstrate the convergence of the perturbation series of the Green's function, it is sufficient to show the convergence of the following series

$$Z = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \frac{d^n}{dE^n} \int d\mu\, \delta(E - H_0)(\lambda S_I)^n \quad . \tag{7}$$

It is worthwhile to stress that the limit $N \to \infty$ is taken after summing the perturbation series. In our previous paper,[2] we took the limit order by order in the series together with the extra condition E = N and obtained the usual

Feynman rules. In general, the infinite volume limit and the summation over the order n of the expansion do not commute. This situation is analogous to that in the standard functional formulation, where an exchange of the functional integration and the summation $e^{-S} = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} S_I^n e^{-S_0}$ is also impossible.

To prove the convergence of the series (7), we scale P_x and ϕ_x as $P_x \to \sqrt{E}\, P_x$ and $\phi_x \to \sqrt{E}\, \phi_x$. The n-th order of the series becomes then

$$A_n \equiv \frac{(-1)^n}{n!} \frac{d^n}{dE^n} E^{N+2n-1} \int d\mu\, \delta(1 - H_0)(\lambda S_I)^n$$

$$= \frac{(-1)^n}{n!} \frac{(N + 2n - 1)!}{(N + n - 1)!} E^{N+n-1} \int d\mu\, \delta(1 - H_0)(\lambda S_I)^n$$

$$< \frac{1}{n!} \frac{(N + 2n - 1)!}{(N + n - 1)!} E^{N+n-1} \int d\mu\, \delta(1 - H_0')(\lambda S_I)^n \quad , \tag{8}$$

where $H_0' \equiv \Delta V \sum_{x=1}^{N} \left(\frac{P_x^2}{2} + \frac{m^2 \phi_x^2}{2} \right)$ and $\Delta V \equiv \frac{V}{N}$.

Using the equality

$$\int d\mu\, \delta(1 - H_0')(\lambda S_I)^n < (\lambda S_I(\phi_x = c))^n \int d\mu\, \delta(1 - H_0') \quad , \tag{9}$$

where $c = \left(\frac{2}{\Delta V\, m^2} \right)^{1/2}$, we find that A_n is less than

$$\frac{(N + 2n - 1)!}{n!(N + n - 1)!} E^{N+n-1}(\lambda S_I(\phi_x = c)^n \int d\mu\ \delta(1 - H_0')$$

$$\xrightarrow[n\to\infty]{} 0\left(\frac{1}{\sqrt{n}}\ (4\lambda\ S_I E)^n\right) \quad . \tag{10}$$

Thus, we have discovered that the perturbation series (7) for given energy E converges absolutely for small enough λ. Summing the series for small λ, we obtain the Green's function by taking the limit $N \to \infty$ afterwards. Before taking this limit, it is necessary to insert for E an explicit value which depends on λ and to make an appropriate analytic continuation in λ. The reason for the former operation is that when we expanded the microcanonical density as in (6), we did not expand the total energy E, which has to be determined from the self-consistency equation (5), but only the interaction part S_I. This is the crux for obtaining a convergent perturbation series.

Next, let us expand the ensemble density with respect to $\lambda^{-1/2}$ (i.e. a strong coupling expansion)

$$\begin{aligned}\int d\mu\ \delta(E - H) &= \int d\mu\ \delta(E - H_0 - \lambda\ S_I)\\ &= \lambda^{-\frac{N}{4}} \int d\mu\ \delta(E - H_I - \lambda^{-\frac{1}{2}} S_0)\\ &= \lambda^{-\frac{N}{4}} \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \frac{d^n}{dE^n} \int d\mu\ \delta(E - H_I)(\lambda^{-\frac{1}{2}}\ S_0)^n\end{aligned} \tag{11}$$

with $H_I \equiv \Delta V \sum_x \left(\frac{p_x^2}{2} + \phi_x^4\right)$,

where we have scaled ϕ_x as $\phi_x \to \lambda^{-\frac{1}{4}}\phi_x$. The n-th order ($=A_n$) of the series can be written as

$$\lambda^{\frac{N}{4}} A_n = \frac{(-1)^n}{n!} \frac{d^n}{dE^n} \int d\mu\ \delta(E - H_I)(\lambda^{-\frac{1}{2}} S_0)^n$$

$$= \frac{(-1)^n}{n!} \frac{d^n}{dE^n} \left(E^{\frac{3}{4}N + \frac{n}{2} - 1}\right) \lambda^{-\frac{n}{2}} \int d\mu\ \delta(1 - H_I) S_0^n \quad . \tag{12}$$

If we note the following equality,

$$S_0 \leqq \Delta V \left\{ \sum_{x,\mu} \frac{\phi^2_{x,\mu} + \phi^2_x}{a^2} + \sum_x \frac{m^2}{2} \phi^2_x \right\} \equiv S_0' \tag{13}$$

it is easy to show that

$$A_n \leqq |A_n| \leqq \frac{\lambda^{\frac{N}{4}}}{n!} \left| \frac{d^n}{dE^n} \left(E^{\frac{3N}{4} + \frac{n}{2} - 1}\right) \right| \lambda^{-\frac{n}{2}} \int d\mu\ \delta(1 - H_I)(S_0')^n$$

$$< \frac{\lambda^{\frac{N}{4}}}{n!} \left| \frac{d^n}{dE^n} \left(E^{\frac{3N}{4} + \frac{n}{2} - 1}\right) \right| \lambda^{-\frac{n}{2}} S_0'(\phi_x = c')^n \int d\mu\ \delta(1 - H_I) \tag{14}$$

where $C' \equiv (\Delta V)^{-\frac{1}{4}} = (\frac{V}{N})^{-\frac{1}{4}}$ and where we have replaced ϕ_x with a possible maximal value of C'. Therefore, we conclude that the series (11) in the strong coupling expansion for given E also converges absolutely for large enough λ.

In the previous discussion we have demonstrated the convergence of the perturbation series for the scalar field theory. For lattice gauge theories, it is also possible to prove the convergence of the strong coupling expansion. In the case of an even number of degrees of freedom (N = even), we obtain a finite series.

Let us write the action of alattice gauge theory as βS, where β is the inverse of the coupling constant squared. The microcanonical partition function is then

$$\int d\mu\ \delta\left(E - \sum_{i=1}^{N} \frac{P_i^2}{2} - \beta S\right) = \sum_{n=1}^{\infty} \frac{(-1)^n}{n!} \frac{d^n}{dE^n}\left(E^{\frac{N}{2}-1}\right) \int d\mu\ \delta\left(1 - \sum_{i=1}^{N} \frac{P_i^2}{2}\right)(\beta S)^n$$

$$\text{with}\quad d\mu \equiv \prod_{i=1}^{N} dP_i \times \prod_{\ell} dU_\ell \quad \text{and} \quad E = \frac{n}{2} + \langle\beta S\rangle \quad . \tag{15}$$

In (15) dU_ℓ is the Haar measure of a link variable and N is the total number of degrees of freedom, which equals (number of lattice points) x $\frac{d(d-1)}{2}$ x (number of gauge degrees of freedom). It is straightforward to show the convergence of the series (15). We find that the effective expansion parameter is $\beta N/2E$, which is less than β, so that a series like (15) in $\beta N/2E$ converges faster than one in β of the standard formulation.

It is interesting that the series in (15) terminates at order $n_c(= \frac{N}{2} - 1)$ if N is even. This implies that the series of the weak coupling expansion, which is obtained by multiplying the series (15) by $\beta^{-(\frac{N}{2}-1)}$, is convergent also.

Finally, we show that under some reasonable assumptions the microcanonical expectation value (2) equals the one derived in the standard functional formulation, which is given by

$$\lim_{N\to\infty} \frac{1}{Z_c} \int \prod_{x=1}^{N} d\phi_x\ \phi(x_1)\ldots\phi(x_n)e^{-S} = G(x_1\ldots x_n) \tag{16}$$

$$\text{with}\quad Z_c \equiv \int \prod_{x=1}^{N} d\phi_x\ e^{-S} \quad .$$

Using the formula, $\frac{1}{2\pi}\int_{-\infty}^{\infty} d\lambda\, e^{i\lambda(E-H)} = \delta(E - H)$ and performing the integrations over P_i, we obtain from eq. (2)

$$\frac{1}{Z_c}\int d\mu\, \delta(E - H)f(\phi) = \frac{1}{Z_c'}\int_{-\infty}^{\infty} d\lambda \prod_{\chi=1}^{N} d\phi_\chi f(\phi)\lambda^{-\frac{N}{2}} e^{i\lambda(E-S)}$$

$$= \frac{1}{Z_c'}\int_{-\infty}^{\infty} d\lambda\, e^{-\frac{N}{2}\log\lambda + i\lambda E + W} G_\lambda(f) \tag{17}$$

with $e^W \equiv \int \prod_{\chi=1}^{N} d\phi_\chi e^{-i\lambda S}$, $G_\lambda(f) \equiv \dfrac{\int \prod_{\chi=1}^{N} d\phi_\chi e^{-i\lambda S} f(\phi)}{e^W}$

and $f(\phi) \equiv \phi(x_1)\ldots\phi(x_n)$.

In (17) we have not specified an irrelevant normalization factor. In order to evaluate the integral over λ as $N \to \infty$, we assume that both of the following limits exist

$$\lim_{N\to\infty} G_\lambda(f) \quad , \quad \lim_{N\to\infty} \frac{\partial}{\partial\lambda} W/N \quad \text{for} \quad \operatorname{Im}\lambda \leqq 0 \quad . \tag{18}$$

Then, by using the steepest descent method as $N \to \infty$ and the relation $E = \frac{N}{2} + \langle S\rangle$, it is easy to derive the standard formula (16). The stationary point required can be found by solving the equation,

$$-\frac{1}{2\lambda} + \frac{iE}{N} + \frac{1}{N}\frac{\partial W}{\partial\lambda} = 0 \quad , \quad E = \frac{N}{2} + \langle S\rangle \quad , \tag{19}$$

where $\langle S\rangle$ is given by (4).

To summarize, we have shown that for a sufficiently small expansion parameter, the perturbation series converges in the microcanonical formulation of the scalar theory and of the gauge theories with an even total number of degrees of freedom. The convergence of the series is only guaranteed for a finite volume. However, if we are able to sum the series and make appropriate analytic continuation in the expansion parameter of the result, we may construct, in principle, all the regularized Green's functions in the infinite volume limit. The reason for being able to obtain non-perturbative effects from perturbation theory is eq. (5), in which we need to estimate non-perturbatively the quantity <S> and then develop the perturbation theory as discussed above.

Acknowledgements

This work was initiated when the author was at the Institute for Nuclear Study of the University of Tokyo. He would like to thank Dr. F.R. Klinkhamer for reading through the manuscript.

This work was supported in part by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

References

1. D.J.E. Callaway and A. Rahman, Phys. Rev. Lett. 49 (1982) 613

2. A. Iwazaki, Phys. Lett. 141B (1984) 342 and INS-Rep 497 (1984)

3. C.S. Lam, Nuovo Cimento 55A (1968) 258

   C. Bender and T.T. Wu, Phys. Rev. Lett. 27 (1971) 461

   L.N. Lipatov, Zh. Eksp. Teor. Fig., Pis'ma Red 25 (1977) 25

   J. Zinn-Justin, Phys. Rep. 49 (1978) 206

4. U.M. Heller and N. Seiberg, Phys. Rev. D27 (1983) 2257

   J. Polonyi and H.W. Wyld, Phys. Rev. Lett. 51 (1983) 2257

   J. Polonyi, H.W. Wyld, J. Kogut, J. Shigemitsu and D.K. Sinclair, Phys. Rev. Lett. 53 (1984) 644 and OSU Preprint, DOE/ER/01545-349

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720