LBL-20301
Preprint c.2

Lawrence Berkeley Laboratory
UNIVERSITY OF CALIFORNIA

RECEIVED
LAWRENCE
BERKELEY LABORATORY
JAN 15 1986
LIBRARY AND
DOCUMENTS SECTION

Submitted to Journal of Physics A

RELATIVISTIC FERMION IN PERIODIC
SQUARE POTENTIAL

B. Banerjee, N.K. Glendenning, and V. Soni

September 1985

NUCLEAR SCIENCE DIVISION

LBL-20301
c.2

Prepared for the U.S. Department of Energy under Contract DE-AC03-76SF00098

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

LBL-20301

Relativistic Fermion in Periodic Square Potential

B. Banerjee, N.K. Glendenning, and V. Soni

Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, CA 94720

*This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

Relativistic Fermion in Periodic Square Potential*

B. Banerjee, N. K. Glendenning, and V. Soni

Nuclear Science Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

ABSTRACT

An analytic solution for a Dirac particle in a one dimensional periodic square potential is obtained. The solution goes over in the non-relativistic limit to the famous Kronig-Penney one, and exhibits similar band structure of the energy spectrum.

*This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U. S. Department of Energy under Contract DE-AC03-76SF00098.

The motion of non-relativistic electrons in a periodic one-dimensional square well potential was solved long ago by Kronig and Penney, as a model of electrons in a crystal [1]. Their solution explicitly exhibits a band structure in the spectrum. There are also several physical systems where relativistic particles feel a periodic potential. The electrons in neutron stars are relativistic and if the neutrons of the dense core form a solid lattice, as has been conjectured, they would experience a periodic potential.

The particular interest that we have in this problem arises from recent interest in soliton models of the nucleon as representing a non-perturbative solution to Q.C.D. . We have studied a model of dense nuclear matter in which the nucleons are represented by quarks that are tightly bound to a topological configuration of meson fields (soliton). The solitons are arranged in a crystaline lattice. In this case the relativistic quarks experience the periodic field of the topological meson configurations. We have posed the question whether matter, described in this way might exhibit a phase transition in which, as in metals, the quarks become liberated from the solitons in which they are normally bound [2].

It is therefor of some interest to solve the relativistic analogue of the Kronig-Penney model.

The Dirac equation, in one space dimension can be written,

$$(\vec{\alpha}\cdot\vec{p} + \beta(m + U) - E)\,\psi(x) = 0 \ , \ \psi = \binom{u}{v} \tag{1}$$

where

$$\alpha = \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \ \beta = \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \tag{2}$$

The resulting linear coupled equations for u and v can be rewritten in terms of one of these, say u, which satisfies (units are $\hbar = c = 1$),

$$u'' + uU' = \Big[(m + U)^2 - E^2 \Big] u \tag{3}$$

For a periodic square potential, U, a discontinuity is produced in the derivative of the solution. For

$$U(x) = \begin{cases} 0 \quad , & |x| \leqslant a \\ V \quad , & a < |x| \leqslant a + b \end{cases} \tag{4}$$

we have

$$U' = -V\Big[\,\delta(x + a) - \delta(x - a)\Big] \tag{5}$$

and hence

$$\begin{aligned} u'' - uV\Big[\,\delta(x + a) - \delta(x - a)\Big] \\ = \Big[\,(m + U)^2 - E^2\Big]u \end{aligned} \tag{6}$$

Integrate this about a small neighbourhood of each discontinuity to find the discontinuity in derivative of the solution,

$$\begin{aligned} u'(-a + \eta) - u'(-a - \eta) &= Vu(-a) \\ u'(a + \eta) - u'(a - \eta) &= -Vu(a) \end{aligned} \tag{7}$$

Now define

$$Q^2 = (m + V)^2 - E^2 \quad , \quad K^2 = E^2 - m^2 > 0 \tag{8}$$

In each of the three regions the solution has the form,

$$\begin{aligned} u_1 &= Ae^{Qx} + Be^{-Qx} \quad , \quad -(a + b) \leqslant x < -a \\ u_2 &= \alpha e^{iKx} + \beta e^{-iKx} \quad , \quad -a \leqslant x < a \\ u_3 &= Ce^{Qx} + De^{-Qx} \quad , \quad a < x \leqslant a + b \end{aligned} \tag{9}$$

The Block theorem [3] can be used to relate the wave function at points separated by an integer number of cell lengths,

$$\begin{aligned} \psi(-a - b) &= \psi(a + b)\exp(\,2i(a + b)k) \\ \psi'(-a - b) &= \psi'(a + b)\exp(\,2i(a + b)k) \end{aligned} \tag{10}$$

which solve to give

$$C/A = e^{-2(a+b)(Q+iK)}\,,\ D/B = e^{2(a+b)(Q-iK)} \tag{11}$$

The continuity of the function, and the specific discontinuity of the derivative derived above yield in particular

$$
\begin{aligned}
u_1(-a-\eta) &= u_2(-a+\eta) \\
u_1'(-a-\eta) + Vu_1(-a+\eta) &= u_2'(-a+\eta) \\
u_3(a+\eta) &= u_2(a-\eta) \\
u_3'(a+\eta) + Vu_3(a+\eta) &= u_2'(a-\eta)
\end{aligned}
\tag{12}
$$

in which the limit $\eta \to 0$ can now be taken. Taking into account (11), these four equations determine all the constants in the solution provided the determinant of the coefficients of the unknowns vanishes

$$
0 = \begin{vmatrix}
1 & 1 & 1 & -1 \\
Q+V & -(Q-V) & -iK & iK \\
e^{-2Qb}e^{-2ik(a+b)} & e^{2Qb}e^{-2ik(a+b)} & -e^{2iKa} & -e^{-2iKa} \\
(Q+V)e^{-2Qb}e^{-2ik(a+b)} & -(Q-V)e^{2Qb}e^{-2ik(a+b)} & -iKe^{2iKa} & iKe^{-2iKa}
\end{vmatrix}
\tag{13}
$$

After some manipulation, this yields

$$
\frac{Q^2 - K^2 + V^2}{2QK} \sinh 2Qb \sin 2Ka + \cosh 2Qb \cos 2Ka = \cos 2k(a+b) \tag{14}
$$

This provides the eigenvalues, E, in terms of the crystal momentum, k. The band structure arises because the left side cannot equal the right except over continuous ranges of energies that are separated by gaps. This is similar to the Kronig - Penney result but differs explicitly in the appearance of V on the left side, and implicitly in the relativistic relations (8). This result embraces the entire range of energies and reduces in the non-relativistic limit to the result of Kronig-Penney.

In Fig 1 we illustrate in a particular example chosen so that the particle is relativistic near and above the top of the potential, how the levels of the isolated potentials develop into bands, which merge for sufficiently close spacing of the wells. The band structure persists into the positive energy spectrum above the top of the potential, with the gaps tending toward zero with increasing energy.

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics, of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

References

1. R. Kronig and W. Penney, Proc. Roy Soc A130 (1931) 451.

2. B. Banerjee, N. K. Glendenning and V. Soni, Phys Lett. 155B (1985) 213.

3. C. Kittel, Introduction to Solid State Physics (John Wiley, New York, 1976)

Figure Captions

Fig. 1. Energy levels and band structure in a one dimensional periodic potential, for a Dirac particle, are shown as a function of separation between the attractive regions. At large separation the levels are degenerate and in the limit coincide with those of the isolated potential of width 2a. In units $\hbar = c = 1$, the mass and potential height are taken equal to 5 fm^{-1} so that the particle is relativistic and at above the potential.



Fig. 1

XCG 851-10

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

LAWRENCE BERKELEY LABORATORY
TECHNICAL INFORMATION DEPARTMENT
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720