Submitted to Chemical Communications

LBL-2593
Preprint c.2

SALTS OF THE HEXAFLUOROBENZENE CATION, $C_6F_6^+$

T. J. Richardson and Neil Bartlett

RECEIVED
LAWRENCE
RADIATION LABORATORY

APR 5 1974

LIBRARY AND
DOCUMENTS SECTION

March 1974

Prepared for the U. S. Atomic Energy Commission
under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy which may be borrowed for two weeks. For a personal retention copy, call Tech. Info. Division, Ext. 5545

LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY

LBL-2593
c.2

25

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Salts of the Hexafluorobenzene Cation, $C_6F_6^+$

By T. J. Richardson and Neil Bartlett

Inorganic Materials Research Division, Lawrence Berkeley Laboratory
and Department of Chemistry; University of California,
Berkeley, California 94720

Summary: Interaction of $O_2^+AsF_6^-$ with hexafluorobenzene in WF_6 solution yields a bright-yellow paramagnetic solid (g = 2.0063), $C_6F_6^+ AsF_6^-$, crystals of which are hexagonal (space group $R\bar{3}$), with $a_o = 10.55$, $c_o = 7.66$ Å, $V = 738.3$ Å^3, $z = 3$, $d_c = 2.53$ g cm^{-3}.

Although the unusually high electron affinities of the heavier transition metal hexafluorides (particularly those of PtF_6 and IrF_6) suggested[1] their utility in synthesising salts of the $C_6F_6^+$ ion (I = 9.97ev)[2], nevertheless attempts by Jha to prepare $C_6F_6^+ PtF_6^-$ resulted in facile fluorination of the C_6F_6 by the PtF_6.[3] More recently we have obtained a bright-orange solid from interaction of IrF_6 with C_6F_6 in tungsten hexafluoride solution, which we believe to be $C_6F_6^+ IrF_6^-$. Unfortunately this compound decomposes quickly on warming towards room temperature. Evidently the hexafluorometallate (v) anions of the noble metals are too labile and too strongly oxidizing for these approaches to $C_6F_6^+$ salt synthesis to be effective.

Since the AsF_6^- ion has considerable kinetic stability it appeared that the $O_2^+ AsF_6^-$ salt, with its potent one-electron oxidizer O_2^+ (E = 12.2ev)[4], would be suitable for $C_6F_6^+$ generation. Tungsten hexafluoride was used as solvent for the hexafluorobenzene and as moderator for the vigorous reaction (fluorocarbon solvents were less satisfactory in the latter function). In a typical experiment C_6F_6 and WF_6 were co-condensed onto O_2ASF_6 powder at -196°, held in a Teflon FEP or quartz reaction vessel. Oxygen was evolved

quantitatively on warming towards 0°, according to the equation:

$$O_2AsF_6 + C_6F_6 \xrightarrow{WF_6} C_6F_6AsF_6 + O_2$$

The yield of $C_6F_6^+ AsF_6^-$ in preparations of this kind is ∿ 40%. The solid product is a bright-yellow powder which decomposes fully in 5-6 hous at ∿ 20° but has been kept for several weeks at -15°.

Infrared specta of the freshly-prepared powder have absorption maxima at 1490(br,s), 1240(w), 1190(m), 1030(s) and 700(vs), compared with 1531(vs), 1368(s) and 1020(vs)cm^{-1} for the gas-phase hexafluorobenzene molecule[5] and 700 cm^{-1} for hexafluoroarsenate ion.[6]

ESR spectra of dilute solutions in HF, as well as of the pure solid, exhibit an unresolved broad symmetric signal at g = 2.0063. Bazhin et al report[7] that solutions of C_6F_6 in superacids give resolved spectra[5] with g = 2.0038 which they attribute to $C_6F_6^+$. Accurate bulk susceptibility measurements have been difficult to obtain because of decomposition in the course of the manipulation of the samples, but the data show that the Curie law is obeyed in the temperature range 3.8 to 71.5°K and that the magnetic moment is in excess of 1.3B.M. (the thermal decomposition products are diamagnetic).

The salt $C_6F_6^+AsF_6^-$ dissolves appreciably in anhydrous HF at 25° and may be recrystallized from it by cooling. The resulting orange crystals, m.p. 69.1° (decomp.), shown to be indentical with the original powder by X-ray

photography, are hexagonal, $R\bar{3}$ (by Weissenberg photography) with $a_o = 10.55(1)$, $c_o = 7.66(1)\mathring{A}$, $V = 738.3\mathring{A}^3$, $z = 3$, $d_c = 2.53$ g cm^{-3}. A complete structure determination is in progress but it is already clear that the $C_6F_6^+$ entity (which is expected to exhibit a Jahn-Teller distortion) must either be disordered in its lattice placement or else possess three-fold axial symmetry.

In accord with its low ionization potential, nitric oxide (I=9.25 ev)[4] interacts quantitatively with $C_6F_6AsF_6$:

$$C_6F_6\,AsF_6 + NO \longrightarrow NOAsF_6 + C_6F_6$$

the C_6F_6 being identified by infrared spectroscopy and the $NO\,AsF_6$ by its characteristic X-ray-powder and Raman spectra.

The dioxygenyl fluoroantimonates ($O_2^+\,SbF_6^-$ and $O_2^+\,Sb_2F_{11}^-$)[8] also oxidize C_6F_6 (in WF_6) to yield $C_6F_6^+$ salts in almost quantitative yield. The former yields a friable solid and the latter a lower melting material. The fluoroantimonates are thermally more stable than the fluoroarsenate but crystalline samples are more difficult to obtain.

This work was supported by The United States Atomic Energy Commission and by The Committee on Research of The University of California, Berkeley.

1. N. Bartlett, Angew. Chem., Int. Ed., 7, 433 (1968).

2. N.B.S. (U.S.A.) Nat. Stand. Ref. Data Ser.-NBS26, 151.

2. N. K. Jha, Ph.D. Thesis, University of British Columbia, 1965.

4. Reference 2, p. 116.

5. D. Steele and D. H. Whiffen, Trans. Farad. Soc. 55, 369.

6. K. Nakamoto, "Infrared Spectra of Inorganic and Coordination Compounds," Second Edition, J. Wiley & Sons, New York, 1970, 122.

7. N. M. Bazhin, Yu. V. Pozdnyakovich, V. D. Shteingarts, and G. G. Yakobson, Izv. Akad. Nauk SSR, Ser. Khim., 10, 2300 (1969).

8. D. E. McKee and Neil Bartlett, Inorg. Chem., 12, 2738 (1973).

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720