

Submitted to Biochemical and Biophysical  
Research Communications

LBL-2646

Preprint *e.2*

FLUORESCENT  $\text{Ca}^{+2}$  ANALOGS IN NERVE:  
RARE-EARTH IONS

Daniel Axelrod and Melvin P. Klein

February 14, 1974

RECEIVED  
LAWRENCE  
RADIATION LABORATORY


MAR. 19. 1974

LIBRARY AND  
DOCUMENTS SECTION

Prepared for the U. S. Atomic Energy Commission  
under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545



LBL-2646  
*e.2*

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

# FLUORESCENT $\text{Ca}^{+2}$ ANALOGS IN NERVE: RARE-EARTH IONS

Daniel Axelrod and Melvin P. Klein

Laboratory of Chemical Biodynamics, Lawrence Berkeley Laboratory,

University of California, Berkeley, California 94720

Received

## SUMMARY

To probe the  $\text{Ca}^{+2}$  ion-nerve membrane interaction, rare-earth ions were used as analogs of  $\text{Ca}^{+2}$  in the bathing solution of garfish olfactory nerve trunk. We observed the chemically sensitive fluorescence of  $\text{Eu}^{+3}$  and  $\text{Tb}^{+3}$  as a measure of rare-earth binding to nerve, both resting and firing, and subjected to treatments with various enzymes, chemical agents, and abnormal salt concentrations. The main results are: 1) rare-earth ions bind to nerve proteins, NADH, and possibly other sites; 2) fractional changes of rare-earth fluorescence synchronous with firing are less than 0.02%; 3) most of the observed binding is a slow, continual uptake of rare-earth ion into the axoplasm, rate-limited by permeation through the axon membrane.

## INTRODUCTION

$\text{Ca}^{+2}$  ions in the external solution of a nerve stabilize the resting state of the axon membrane and play an essential role in excitability (1). We substituted the fluorescent rare-earth ions,  $\text{Eu}^{3+}$  and  $\text{Tb}^{3+}$ , for  $\text{Ca}^{+2}$ , and observed their characteristically enhanced fluorescence intensities upon chemical binding under varying conditions of nerve firing, solution ion concentrations, enzyme treatments, and chemical agents, as measures of the quantity of rare-earth ions bound to the nerve.

Chlortetracycline has been used previously as a fluorescent probe for  $\text{Ca}^{+2}$  bound to nerve, with results of ambiguous interpretation (2). Rare-earth ions have been used as fluorescent probes of metal binding sites in the protein transferrin (3).

## MATERIALS AND METHODS

The nerve tissue used for this study was the garfish olfactory nerve trunk, which has a uniquely high ratio of axon membrane surface area to volume and to Schwann cell and connective tissue area (4). The dissections were performed as described by Easton (4). All experiments were performed at room temperature. The compositions of the various Ringer solutions used are given in Table 1. The enzymes were obtained commercially and used

without further purification.

All the fluorescence emission and excitation spectra (uncorrected) were recorded with a Perkin-Elmer MPF-2a spectrofluorimeter with a spectral resolution of 5 nm. The various spectra of nerve were recorded by pushing a 4 cm length of garfish nerve trunk into a 2 x 2 mm quartz fluorescence cuvette filled with Ringer solution containing the appropriate ions.

A separate apparatus was constructed for experiments examining fluorescence changes. The light from a 900 W Xenon lamp was passed through a water filter, interference filters, and a grating monochromator, yielding an excitation beam of 10 nm spectral bandwidth. A glass plate deflected part of the exciting light directly to a reference photomultiplier, used to monitor the intensity in the Xenon arc. The width of the main beam impinging on the nerve was 0.5 mm after focusing and passage through the quartz side window of the nerve holder. Fluorescent light from the nerve was viewed at 90° through a microscope and interference filters by a photomultiplier mounted above the microscope. The optical set-up was contained completely in a light-tight box during experiments.

The photomultiplier outputs were connected to opposite inputs of a differential amplifier, so that lamp intensity fluctuations would not affect the amplifier output. For the experiments on firing nerves, the amplifier output was filtered to pass 1 Hz to 1 kHz and connected directly to the input of a Computer of Average Transients (CAT) in order to signal average the photomultiplier outputs during several thousand synchronous stimulations of the nerve. For the experiments on resting nerves undergoing changes of bathing solution, the differential amplifier output was DC-coupled with a 12-sec time constant into a strip chart recorder.

Solutions flowed by gravity into the nerve holder and could be switched rapidly without interrupting the flow or disturbing the nerve or optics. The nerve holder itself contained quartz glass windows for passage of the exciting and fluorescent light, provision for the flow of solution into and out of the chamber, plastic supports to prevent nerve sagging, and two pairs of external platinum electrodes for nerve stimulation and recording.

## RESULTS

Rare-earths as Physiological Substitutes for  $\text{Ca}^{+2}$ . Previous work indicates that several polyvalent cations ( $\text{La}^{+3}$ ,  $\text{Ba}^{+2}$ ,  $\text{Ni}^{+2}$ ,  $\text{Co}^{+2}$ ,  $\text{Cd}^{+2}$ ) can substitute for  $\text{Ca}^{+2}$  in the Ringer solution and still maintain nerve excitability (5-7). These polyvalent cations have a physiological effect somewhat similar to that of high  $[\text{Ca}^{+2}]$  (8). Because of the fairly non-specific nature of successful  $\text{Ca}^{+2}$  replacements and the chemical homology of the lanthanides, one would anticipate that any of the trivalent rare-earth ions would successfully substitute for  $\text{Ca}^{+2}$ . This substitution was tested on garfish olfactory nerve by bathing each of several nerves in solutions

containing either one type of polyvalent ion or none at all. The ions used were  $\text{Ca}^{+2}$ ,  $\text{Ni}^{+2}$ ,  $\text{Mn}^{+2}$ , and  $\text{Eu}^{+3}$ . The excitability of the nerves was checked frequently over a period of hours by monitoring the relative persistence of the compound action potential (Fig. 1). The results for Ca-, Eu-, and Ca-free-Ringers indicate that: 1) Garfish nerves maintain excitability in Ca- and Eu-Ringers longer than in Ca-free-Ringer; 2) bathing a nerve in Ca- or Eu-Ringer previously made inexcitable by Ca-free-Ringer restores some excitability. On the basis of such evidence, we proceeded with the use of  $\text{Eu}^{+3}$  and a homologous rare-earth ion,  $\text{Tb}^{+3}$ , as physiological substitutes for  $\text{Ca}^{+2}$ .

Rare-earth Fluorescence Spectra in Nerve. The fluorescence intensity of most of the emission bands of both Eu and Tb aqueous solutions increases upon chemical binding of the ions to strong ligands (9). Comparison of the emission spectra of aqueous  $\text{EuCl}_3$  vs.  $\text{Eu-EDTA Cl}_3$  (Fig. 2), shows that upon  $\text{Eu}^{+3}$  complexation, all emission bands are greatly enhanced, but the relative intensities of the 590 nm and 614 nm peaks become reversed. This reversal can be taken as a definite indication that at least some  $\text{Eu}^{+3}$  in a sample is chemically bound to ligands. The fluorescence spectrum of nerve bathed at 0°C for 24 hr in Eu-Ringer (Fig. 3) also shows this intensity reversal, indicating that a considerable portion of the Eu associated with the nerve was chemically bound. The background fluorescence is probably from oxidized flavins (10).

The fluorescence excitation spectrum of Eu-nerve was obtained point-by-point by measuring the height of the  $\text{Eu}^{+3}$  614 nm peak above the background as a function of excitation wavelength. The resulting (uncorrected) spectrum shows a broad band in the 320-380 nm region not present in the aqueous Eu excitation spectrum (Fig. 3). This band must result from energy transfer from an organic species near the bound  $\text{Eu}^{+3}$  which absorbs in this wavelength region, probably NADH. The excitation spectrum of nerve bathed in

Tb-Ringer using the emission line at 543 nm (Fig. 4) shows a strong band in the 290-300 nm region not present in the aqueous  $Tb^{+3}$  excitation spectrum. This band indicates energy transfer from, and probable  $Tb^{+3}$  binding to, proteins. Since  $Eu^{+3}$  and  $Tb^{+3}$  are similar in chemistry, we conclude that both probably are bound to NADH and proteins. In vitro 1:3 mixtures of  $Eu^{+3}$  and NADH at pH 7.8 were found to exhibit a fluorescence emission characteristic of bound  $Eu^{+3}$  and an excitation spectrum indicating energy transfer from NADH to  $Eu^{+3}$ .

Transient Changes of Eu-Nerve Fluorescence. Fractional transient changes of  $Eu^{+3}$  ( $\lambda=614$  nm) or  $Tb^{+3}$  ( $\lambda=543$  nm) fluorescence, synchronous with the passage of an action potential, were less than 0.02%, the noise limit. This result was obtained irrespective of (1) excitation wavelength or (2) the time the nerve had been bathed in Eu-Ringer before the start of the experiment, from a few minutes to a day. Similarly, transient changes of intrinsic fluorescence from proteins, NADH, and oxidized flavins, both for total and polarized fluorescence, were less than 0.01%. Each of these experiments was tried on at least three garfish nerves.

Fluorescence of Eu-Resting Nerve. The  $\lambda=614$  nm  $Eu^{+3}$  fluorescence of a resting nerve was monitored as a function of time. The bathing solution flowing through the nerve holder could be switched between Ca-Ringer and Eu-Ringer at any time. The  $Eu^{+3}$  fluorescence, presumably proportional to  $Eu^{+3}$  uptake, rose at a fairly linear rate, at least for the first 7 hr. Switching the solution back to Ca-Ringer caused no decrease in fluorescence; i.e., the  $Eu^{+3}$  uptake is irreversible. With Eu-Ringers containing only 0.01 mM  $Eu^{+3}$ , the rate of uptake was less, but still linear, even at early times. Differences in the rate of uptake between a resting nerve and the same nerve stimulated at the rate of 1 pulse/sec were immeasurably small. The time required for uptake of the membrane-impermeable dye, trypan blue, was

measured to be <1 hr, suggesting that the gradual  $\text{Eu}^{+3}$  uptake was not limited by diffusion into the extracellular spaces between the axons.

Since a high  $[\text{K}^+]$ -induced depolarization might change the  $\text{Eu}^{+3}$  binding state, we observed  $\text{Eu}^{+3}$  fluorescence as the solutions were switched between low and high  $[\text{K}^+]$  Eu-Ringer. A small change was observed, but it might result from a simultaneously observed light scattering change.

Eu-Ringer solutions containing either of two proteases, trypsin (1 mg/ml) or pronase (1 mg/ml), or the enzyme neuraminidase (0.1 mg/ml) affected Eu binding to nerve by less than 0.5%. However, treatment by either phospholipase-c (0.2 mg/ml) or dimethylsulfoxide (which increases membrane permeability) led to a 2- to 4-fold increase in the rate of Eu uptake. The dimethylsulfoxide effect was reversible; the phospholipase-c effect was irreversible.

#### DISCUSSION

The results of this work are: (1) continual, slow uptake of  $\text{Eu}^{+3}$  by nerve; (2) enhancement of the rate of uptake after membrane alteration by phospholipase-c and dimethylsulfoxide; and (3) spectroscopic evidence of rare-earth binding to protein and NADH in the nerve. These are all consistent with the interpretation that the bulk of  $\text{Eu}^{+3}$  fluorescence results from Eu binding in the axoplasm, possibly to the mitochondria and as organic complex precipitates, rate limited by permeation through the axon membrane. Ca is known to accumulate in mitochondria (11). The small fraction of Eu possibly bound to the axon membrane may undergo binding changes during nerve stimulation, but the background of axoplasmic  $\text{Eu}^{+3}$  and intrinsic fluorescence interfered with optical detection of these changes, if any.

The technique of replacement of  $\text{Ca}^{+2}$  with a fluorescent rare-earth analog might be especially suited to probing  $\text{Ca}^{+2}$  binding sites in internally perfused squid axons, where the axoplasmic composition can be controlled, in systems in which Ca involvement is greater, as in muscle, and in more homogeneous in vitro biochemical systems.

REFERENCES

1. Tasaki, I. (1968) Nerve Excitation--A Macromolecular Approach, pp. 112-116, Charles C. Thomas, Springfield.
2. Hallett, M., Schneider, A., and Carbone, E. (1972) J. Membrane Biol. 10, 31-44.
3. Luk, C. K. (1971) Biochemistry 10, 2838-2843.
4. Easton, D. (1971) Science 172, 952-955.
5. Blaustein, M. P., and Goldman, D. E. (1968) J. Gen. Physiol. 51, 279-291.
6. Takata, M., Pickard, W. E., Lettvin, J. Y., and Moore, J. W. (1966) J. Gen. Physiol. 50, 461-471.
7. Khodorov, B. I., and Peganov, E. M. (1969) Biofizika 14, 474-484.
8. Frankenhauser, G., and Hodgkin, A. L. (1957) J. Physiol. 137, 218-244.
9. Sinha, S. (1967) Europium, pp. 100-134, Springer-Verlag, New York.
10. Radda, G. K., and Dodd, G. H. (1968) In: Luminescence in Chemistry, (Bowen, E. J., ed.), pp. 192-221, Van Nostrand, London.
11. Reynafarje, B., and Lehninger, A. L. (1969) J. Biol. Chem. 244, 584-593.

Table 1. Compositions of various garfish Ringer solutions used  
(Concentrations in mM/liter)


Ringer name Salt	Ca	Ca-free	Eu	Tb	High[K <sup>+</sup> ]Eu	Low Eu
NaCl	204	204	204	204	3.5	204
KCl	3.5	3.5	3.5	3.5	204	3.5
CaCl <sub>2</sub>	3.5	-	-	-	-	-
EuCl <sub>3</sub>	-	-	0.5	-	0.5	0.05
TbCl <sub>3</sub>	-	-	-	0.5	-	-
Glucose	24	24	24	24	24	24
pH (Hepes Buffer)	6.9	6.9	6.7	6.7	6.7	6.9

Fig. 1. Each trace is the action potential of the nerves bathed in the indicated solutions and elicited at the times (in hr) shown at the beginning of the trace. The duration of each trace is 500 ms. For example, the three traces on line IV show the action potential of a nerve: (1) at  $t=0$  hr, immediately after placement in Ca-free-Ringer; (2) at  $t=16$  hr, still in Ca-free-Ringer; and (3) at  $t=24$  hr, after bathing in Eu-Ringer for the last 6 hr.

Fig. 2. Fluorescence emission spectra of  $\text{EuCl}_3$  (25 mM) aqueous solution (dashed line) vs. Eu (25 mM) - EDTA (50 mM) solution (solid line). Resolution = 2.0 nm,  $\lambda=614$  nm,  $\lambda_a=393$  nm.

Fig. 3. (a) Fluorescence emission spectra of  $\text{EuCl}_3$  aqueous solution vs. Eu-nerve.  $\lambda_a=393$  nm. Resolution = 5.0 nm. (b) Fluorescence excitation spectra (uncorrected) of  $\text{EuCl}_3$  aqueous solution vs. Eu-nerve.  $\lambda=614$  nm. Resolution = 5.0 nm.


Fig. 4. Fluorescence excitation spectra (uncorrected) of  $\text{TbCl}_3$  aqueous solution vs. Tb-nerve.  $\lambda=543$  nm. Resolution = 5.0 nm.



Garfish nerve action potentials

Fig. 1

XBL738-3786



XBL738 - 3787

Fig. 2





Fig. 3



XBL738-3789

Fig. 4

LEGAL NOTICE

*This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720